

*ENGINEERING  
SOILS MAP OF FAYETTE  
COUNTY, INDIANA*

*JUNE 1967  
NO. 17*

*Joint  
Highway  
Research  
Project*

*by*  
**A.K.TURNER**

**PURDUE UNIVERSITY  
LAFAYETTE INDIANA**

Interim Report  
ENGINEERING SOILS MAP  
OF FAYETTE COUNTY INDIANA

TO: Dr. G. A. Leonards, Director  
Joint Highway Research Project

June 20, 1967

File: 1-5-2-43

FROM: H. L. Michael, Associate Director  
Joint Highway Research Project

Project: C-36-51B

The attached report, entitled "Engineering Soils Map of Fayette County, Indiana," completes a portion of the project concerned with development of county engineering soils maps of the State of Indiana. This is the 43rd report in the series. The report was prepared by A. K. Turner, Research Instructor, Joint Highway Research Project.

The soils mapping of Fayette County was performed primarily by using the soil survey map sheets published by the Soil Conservation Service, United States Department of Agriculture in the soil survey of Fayette and Union Counties. Airphoto interpretation techniques were used to supplement the pedological data. The resulting Engineering Soils Map is presented as a blackline print.

Respectfully submitted,

*Harold L. Michael/jgs*  
Harold L. Michael  
Associate Director

HLM:ss

Attachments: One

Copy: F. L. Ashbaucher  
W. L. Dolch  
W. H. Goetz  
W. L. Grecco  
G. K. Hallock  
R. H. Harrell

V. E. Harvey  
J. F. McLaughlin  
F. B. Mendenhall  
R. D. Miles  
J. C. Oppenlander  
C. F. Scholer

M. B. Scott  
W. T. Spencer  
F. W. Stubbs  
H. R. J. Walsh  
K. B. Woods  
E. J. Yoder

Interim Report  
ENGINEERING SOILS MAP  
OF INDIANA

TO: Dr. G. A. Leonard, Director  
Joint Highway Research Project

FROM: H. L. Michael, Associate Director  
Joint Highway Research Project

File: 1-2-2-43  
Project: C-36-718  
June 20, 1961

The attached report, entitled "Engineering Soils Map of Fayette County, Indiana," constitutes a portion of the project concerned with development of county engineering soils maps of the State of Indiana. This is the first report in the series. The report was prepared by A. K. Turner, Research Instructor, Joint Highway Research Project.

The soils mapping of Fayette County was performed primarily by using the soil survey map sheets published by the Soil Conservation Service, United States Department of Agriculture in the soil survey of Fayette and Union Counties. Aerialo interpretation techniques were used to supplement the pedological data. The resulting Engineering Soils Map is presented as a blackline print.

Digitized by the Internet Archive  
in 2011 with funding from

LYRASIS members and Sloan Foundation; Indiana Department of Transportation

HLM:as  
Attachments: One

Copy: F. L. Ashbaugh  
W. E. Folsch  
W. H. Goetz  
W. L. Grisco  
O. K. Hallack  
H. H. Harrell

V. E. Harvey  
J. F. McLaughlin  
V. E. Mendenhall  
R. D. Miles  
J. C. Ogonlander  
C. P. Scholer

M. E. Scott  
W. T. Spencer  
E. W. Stubbins  
H. R. J. Wain  
K. B. Woods  
E. J. Yoder

Interim Report

ENGINEERING SOILS MAP  
OF  
PAYETTE COUNTY, INDIANA

by  
A. K. Turner  
Research Instructor

Joint Highway Research Project

Project: C-36-51B

File: 1-5-2-43

Prepared as Part of an Investigation

Conducted by

Joint Highway Research Project  
Engineering Experiment Station  
Purdue University

in cooperation with

Bureau of Public Roads

U. S. Department of Commerce

The opinions, findings and conclusions expressed in this  
publication are those of the authors and not necessarily  
those of the Bureau of Public Roads.

Not Released for Publication

Subject to Change

Not Reviewed By  
Indiana State Highway Commission  
or the  
Bureau of Public Roads

Purdue University  
Lafayette, Indiana

June 20, 1967

ENGINEERING SOILS MAP  
OF  
FAYETTE COUNTY, INDIANA

by  
A. K. Turner

INTRODUCTION

Development of an engineering soils map of Fayette County was the primary goal of this project. The map is appended to this report; the report supplements the engineering soils map information.

The detailed pedological soils maps published in the 1960 Soil Survey of Fayette and Union Counties by the United States Department of Agriculture (7) were the single most important source of data used in the project. These agricultural soils map sheets, at a scale of 1:15,840, were assembled to form a mosaic map of Fayette County. Careful study of the soil series descriptions enabled the grouping of the series into appropriate landform and parent material categories. Preliminary landform and parent material boundaries were then delineated on the mosaic-map.

Routine airphoto interpretation techniques supplemented the pedological data. Aerial photographs were examined and the preliminary boundaries checked and modified, if necessary, to produce final landform and parent-material boundaries. The photographs were contact prints at an approximate scale of 1:20,000. Date of photography was 1940.

Published geologic reports were studied to verify and amplify the soils information. Most important of these were the reports of Dr. Ansel M. Gooding, Department of Geology, Earlham College, Richmond, Indiana (2,3,4). He has made detailed studies of the glacial deposits in this area. His report on the terraces along the Whitewater River (2) contains much



detailed information beyond the scope of this report.

The final landform and parent-material boundaries were graphically reduced to produce the engineering soils map. Symbols were used to delineate the parent materials (grouped according to landform and origin). Textural symbols were then superimposed to indicate the relative compositions of the parent materials. The map also includes a set of soil profiles which indicate the general soil profiles of topographically high and low sites in each parent material area. Each profile shows the general range in depth and texture (ISHC textures) of each soil horizon. Because of the obviously bad construction characteristics of highly organic top-soils, these materials were carefully mapped. However not all of the identifiable areas of these materials were large enough to be shown on the relatively small scale engineering soils map.

#### DESCRIPTION OF THE AREA

##### General Nature of the County

Fayette County is located in east-central Indiana, separated from the Indiana-Ohio State-line by Union County (Figure 1). It has a maximum north-south length of 18 miles, and a width of 14 miles. Total area is 215 square miles (6). Connersville, the county seat, is located in the east-central part of the county about 55 miles east-southeast of Indianapolis, and reports a 1960 population of 17,698 out of a total county population of 24,454 (8). Several small to medium sized industries are located in Connersville, however about 90 percent of Fayette County is farmed (7). Agriculture is mainly concerned with the raising of livestock, chiefly hogs. Thus considerable cleared land, and much woodland, is devoted to pasturing; while forage crops, especially corn, hay, and wheat, are the main crops (7).



**FIGURE 1**  
**LOCATION MAP OF FAYETTE COUNTY**

### Climate

Table 1, derived from the Agricultural Soil Report summarizes temperature and precipitation data obtained at Rushville, Indiana, and believed typical of Fayette County conditions (7).

Fayette County has a continental type of climate with erratic temperature changes within and between seasons. The winters are moderately cold; the summers warm and humid. Analysis of the mean monthly temperatures shows the county to have a freezing index of 192 degree days. Yoder (11) shows that this might indicate a frost penetration of 20 inches in a well-drained non-frost-susceptible base course. The winter season is particularly marked by rapid temperature changes. Commonly periods of two or three days of subzero weather are followed by short periods of warm weather. As a consequence damage from freezing and thawing of highway subgrades can be expected.

Rainfall varies from season to season. Heaviest rains occur in the spring; flooding of the Whitewater River is common.

### Physiography, Topography, and Drainage

Fayette County lies within the Till Plains Section of the Central Lowland Province (1, 10). However, since the glacial drift is very thin over most of the southern and central parts of the county, Malott (5) has included essentially all of the county within the northern portion of the Dearborn Upland physiographic region of the State.

Elevations in the county range from a low of about 730 feet, where the West Fork Whitewater River leaves the county, to a high slightly in excess of 1100 feet, found in the western and northwestern parts of the county. Figure 2B gives a general picture of the topography of the county. The surface is that of a glacial plain conforming somewhat to the irregularities of the underlying bedrock surface. The uplands have been much dissected



TABLE 1

Temperature and Precipitation Data for Fayette County<sup>\*</sup>

Month	Temperature (°F)			Precipitation (inches)		
	Mean	Extreme High	Extreme Low	Mean	Extreme Dry	Extreme Wet
Jan.	28.7	70	-26	3.39	0.71	12.06
Feb.	29.9	71	-25	2.83	0.32	8.31
Mar.	39.9	86	-10	3.96	0.04	12.08
April	50.4	89	14	3.69	0.90	9.11
May	61.6	96	26	4.10	0.53	8.46
June	70.2	101	35	3.93	0.71	8.45
July	74.2	108	40	3.33	0.20	7.96
Aug.	71.6	103	36	3.15	0.29	9.64
Sept.	61.3	101	22	3.38	0.17	9.36
Oct.	54.4	90	12	2.83	0.23	9.11
Nov.	41.3	83	-10	3.17	0.37	9.25
Dec.	31.0	68	-21	2.97	0.48	6.14
Annual	51.5	108	-26	40.73	0.04	12.08
Number of years of records	67	75	75	77	77	77

\*

This data obtained from Agricultural Soil Survey of Fayette and Union County by USDA. The recording station was at Rushville Indiana, in adjacent Rush County, since 1948; prior to 1948 the station was at Mauzy Indiana. Rushville station location is Lat. 39° 36'N.; Long. 85° 27'W. ground elev. = 955 feet.

adjacent to the valleys of the West Fork Whitewater River and its tributaries where local relief is about 250 feet. The valley is fairly wide with extensive multiple terraces in the northern part of the county; it narrows in the southern part.

With the exception of the west-central and northwestern parts of Fayette County, which lie within the East Fork White River drainage basin, the county lies within the Whitewater River drainage basin. The West Fork Whitewater River is the trunk stream for most of the county. Major tributaries are Williams, Fall, Garrison, Village, Wilson, and Bear Creeks. Small areas along the eastern county boundary drain into the East Fork Whitewater River. Figure 2A shows the drainage system within the county.

## Geology

### Bedrock Geology

Bedrock underlying the western third of the county is of Silurian age; the remainder is of Ordovician age. Limestone and interbedded limestone and shale are the common rock types. Rock outcrops are restricted to stream beds and valley walls. They become larger and more numerous in the southern part of the county where many creek-beds are littered with small blocks of flaggy limestone. Most streams in the southern part of the county are rock-controlled.


### Glacial Geology

Glacial deposits belonging to the older Illinoian and the younger Wisconsin ice sheets are found in Fayette County. Each of these ice sheets advanced and retreated several times. It is believed that the Illinoian ice sheet advanced and retreated over the county at least three times (3). Consequently three Illinoian tills, separated by outwash

FIGURE 2A

# TOPOGRAPHIC MAP FAYETTE COUNTY INDIANA

HENRY CO.



WAYNE CO.



FRANKLIN CO.






gravels and sands or eolian silts, have been identified in the area (3, 4).

At least three different Wisconsin advances affected parts of Fayette County. The first advance covered the northeast half of the county, as shown in Figure 3. A distinctive red-brown "Whitewater till" has been found in deep excavations in Fayette and adjacent counties (3, 4). Following retreat of this ice-sheet, a new advance occurred. "Shelbyville" drift deposited from this glacier covers most of the county, with the exception of a south-central area (see Figure 3). A later, much weaker, readvance barely extended into the northwest part of the county. This "Champaign" drift forms a weakly developed ridge moraine and is the youngest glacial deposit in the county.







A complex series of terraces is found along the West Fork Whitewater River and some of its tributaries. Gooding (2) has identified six levels and related these to the various glacial advances. For engineering purposes a two-fold division, into "high" and "low" terraces, appears adequate. The higher terraces are older than the lower terraces, and therefore have different origins. It is thus not surprising that the terrace levels have somewhat different parent materials-the upper terraces are composed of a silt cover overlying sands and gravels; the lower terraces are generally more variable, their compositions ranging from sand and gravel to sand and silt. The lower terraces lack a consistent silt cover common to the upper terraces.

Wisconsin materials exposed at the surface are of either "Shelbyville" or "Champaign" age. The older Shelbyville deposits are more dissected than the younger Champaign areas. A "loess blanket," one to five feet thick, covers the Shelbyville tills, but is thinner or absent on the Champaign till.





### Legend

- |   |   |   |   |
|---|---|---|---|
|  | "Champaign" Drift Boundary                                |   | Highway                                   |
|  | "Shelbyville" Drift Boundary                              |   | County Line                               |
|  | Buried Early Wisconsin Till Boundary ("Whitewater Drift") |  | Location of Sections Described by Gooding |

**FIGURE 3**

## **MAP SHOWING LOCATION OF WISCONSIN DRIFT BOUNDARIES IN FAYETTE AND UNION COUNTIES**

(after A.M. Gooding 1961, 1963)

The accompanying engineering soils map differentiates areas where this loess cover is approximately three or more feet thick. The younger Champaign drift forms a weak ridge moraine extending across the northwest corner of the county. Gently rolling topography, generally higher elevations, and an occasional closed depression serves to separate this area from the flatter (except where later erosion has occurred) Shelbyville till plain.


Older geological reports (5) suggest the presence of two north-south trending ridge moraines, one on each side of the West Fork Whitewater River valley. Airphoto interpretation and topographic map study tends to discount the presence of these "ridge moraines." The areas on each side of the valley are much dissected by stream erosion. However no true closed basins exist and the maximum elevations are lower than the uplands further from the valley. The present study indicates that, for engineering purposes at least, these areas are best classified as till plains.

Several eskers and kames occur on the east side of the West Fork valley south-east of Connersville. Two valleys, which at one time acted as spillways carrying glacial meltwaters from Union County to the east, enter the West Fork valley at this point. One spillway trends northeast toward Lyonsville; the other trends east toward Alquina. The eskers and kames are apparently closely related to these spillways since they are rare in other parts of the county.

## LANDFORMS AND ENGINEERING SOIL AREAS

### Landforms

Fayette County contains a variety of landforms. Five major categories are easily recognized. They are as follows:- 1) Ridge Moraine of Late Wisconsin age, 2) slightly dissected Till Plains of Early Wisconsin age, 3) dissected Illinoian Till Plains overlying bedrock at shallow depths,



**FIGURE 4**  
**LANDFORMS OF FAYETTE COUNTY**

**Legend**

- 1 - Wisconsin Ridge Moraine
- 2 - Wisconsin Ground Moraine (dissected)
- 3 - Terraces
- 4 - Illinoian Ground Moraine
- 5 - Floodplain

(after Fig. 8, Soil Survey of  
 Fayette and Union Counties)

4) River Terraces, and 5) river Flood Plains. Figure 4 shows in an idealized fashion the general relationships of these landforms. There are also several minor landforms, including eskers, kames, small outwash plains, and small loess deposits.

The six different terrace levels described by Gooding (2) have here been grouped into two types:- high terraces which are generally 20-35 feet above the flood plain and consequently are well drained and low terraces which are generally only 5-15 feet above the flood plain and are usually poorly drained. Also, the composition of the terraces materials is generally different; the high terraces normally consist of gravel and sand, overlain by silty sands, while the low terraces are more variable, varying in composition from sand and gravel to sand and silt.

#### Engineering Soil Areas

The soils of Fayette County can be divided into four major groups which reflect their origins:- (a) Glacial deposits, (b) Fluvial deposits, (c) Eolian deposits, and (d) Miscellaneous deposits. These can be further subdivided by landform and parent materials into a number of distinctive units as shown on the engineering soils map and as listed in Table 2. Table 2 also shows the relationships between these landform-parent material subdivisions and the soil series names employed by the pedologists in mapping the soils of Fayette County (7).

#### A) GLACIAL DEPOSITS

Upland areas throughout the county are mantled with a variety of glacial deposits. In the south-central part of the county a thin layer of Illinoian ground moraine overlies limestone and shale bedrock. Most of the remaining upland areas are covered with Wisconsin ground moraine overlain by a thin wind-blown silt layer. A low ridge moraine formed by

TABLE 2

Summary of Landforms and Materials  
in Fayette County

ORIGIN	LANDFORM	PARENT MATERIAL	SOIL SERIES
GLACIAL DEPOSITS	Ridge Moraine (Young Wisconsin)	Silty-clay till with sand lenses	Miami, Crosby
	Wisconsin Ground Moraine (Older Wisconsin)	Clayey-silt overlying silty-clay till	Fincastle, Xenia, Russell, Miami, Hennepin, Celina
	Illinoian Ground Moraine	Clayey-silt till (rock often at shallow depths)	Avonburg, Cincinnati, Hennepin
	Eskers and Kames	Outwash, sands and gravels	Fox (kame phase), Rodman
FLUVIAL DEPOSITS	Terraces a) High Level	Silt over outwash; sand and gravel	Ockley (With minor Fox), Rodman on scarps
	b) Low Level	Outwash; sand and gravel or silt and sand, variable	Fox, Rodman on scarps; (minor Homer, Westland, Martinsville, Whitaker)
	Flood Plains	Silt and sand, alluvium	Genesee, Eel, Nineveh, Ross, Shoals, Sloan
EOLIAN DEPOSITS	Plain	Silt, loess	Manlove, Birkbeck, Reesville
MISCELLANEOUS DEPOSITS	Plateau	Interbedded limestone and shale on the surface or at shallow depths	Fairmount, Milton, Wyun
	Depression	Cumuloose drift	Cope, Brookston, Kokomo



a readvance of Wisconsin ice extends across the northwest corner of the county. Some kames and eskers are associated with the other Wisconsin deposits.

#### Ridge Moraine - Silty Clay Texture

A low ridge moraine, part of the Shelbyville morainic system, extends east-west across the northwestern part of the county. Local relief in this moraine is not great, rarely exceeding 20 feet except near a creek in the northeast where local erosion has occurred.

The soils are dominantly silty clays. The A-horizon, is usually a silty clay loam; The B- and C- horizons are either a clay, or silty clay.

#### Ground Moraine - Silty Texture

The ground moraines lie to the south of, and are older than, the ridge moraine in the northwest corner of the county. Most of these soils have some wind-blown silt on the surface. This serves to greatly increase the silty texture of the near surface horizons, and in places reaches five feet thick. Areas where this silt deposit is over three feet thick have been mapped as separate loess areas.

The silt cover has tended to smooth the rolling topography so that uneroded areas are quite level. However this area has been considerably dissected by the many small tributaries of the Whitewater River. This dissection increases greatly near the Whitewater River and its major tributaries so that very little level topography remains.

The upper portions of soil profiles are very silty; the lower parts of the profiles are silty clays or clays. The A-horizon may be a foot thick and is usually a silty loam or silty clay loam. The B-horizon is normally a silty clay and the C-horizon a clay or silty clay.

### Illinoian Ground Moraine - Silty Texture

Illinoian Ground Moraine is found in the south-central part of Fayette County. On level upland areas 10 to 60 inches of wind-blown silt overlie the Illinoian till which is more clayey. Extensive areas of steep slopes occur due to dissection by the many creeks and gullies. In these areas very thin soil profiles are found overlying interbedded limestone - shale bedrock.

In upland areas the A-horizon is about 12 inches thick and is classified as a silty loam. The B-horizon extends to depths of 60 to 78 inches and is generally a silty clay or clay. The C-horizon is also a silty clay or clay and normally contains only a very few small pebbles.

Where erosion has occurred the soil profile is generally less than a foot deep with a silty textured A-horizon a few inches thick, and a B-horizon of silty-clay extending down to about 12 inches. Where rock is very close to the surface flat slabs of limestone are common throughout the soil profile.

### Eskers and Kames - Sandy and Gravelly Texture

Several kames and short esker-like ridges are found in Fayette County. A concentration of these deposits is found on the east side of the Whitewater valley, southeast of Connersville, and extends in a discontinuous fashion eastward towards Alquina.

The depths and compositions of the various soil profile layers are quite variable. Frequently the A-horizon may be a silty, sandy, or even clayey loam. The depth of the A-horizon ranges from a few inches to perhaps a foot. The underlying B-horizon ranges in composition from silty clay to sandy clay or gravelly clay and may extend to depths of about three feet. The underlying parent material is sand and gravel, sometimes with small amounts of intermixed or interbedded silt and clay.

## B) FLUVIAL DEPOSITS

Extensive fluvial deposits are found along the West Fork Whitewater River and along the lower portions of its many small tributaries. Small outwash plains and terraces are found along the abandoned glacial sluiceways in the eastern part of the county. The West Fork Whitewater River exhibits an extensive flood plain, or alluvial plain, and the many creeks exhibit proportionately broad flood plains.

### Terraces - Sandy and Gravelly Texture

The terraces along the Whitewater River and its tributaries are generally composed of sands and gravels. These terraces are most extensive in the vicinity of Connersville, although examples of these terraces can be found in other parts of the county.

Two terrace levels can be distinguished. The upper level terraces occur from 20 to 35 feet above the flood plain. These terraces normally have a silty sand cover about three feet thick overlying the sand and gravel. Accordingly the A-horizon can be classified as a silty loam and is often a foot thick. The B-horizon gradually changes from a silty clay to a sandy or gravelly clay with depth. Below about five feet sand and gravel (C-horizon) is encountered.

The lower terraces occur from 5 to 15 feet above the flood plain and normally lack the silt cover so common on the upper terraces. The A-horizon is usually a foot thick and is generally a loam. The B-horizon rapidly grades downward into a sandy or gravelly clay and rarely extends below three feet. The underlying parent material is stratified sand and gravel, or sand and silt, and is quite variable.

### Alluvial Plains

The alluvial plains shown on the map include the flood plains of the various rivers and creeks and so can be considered as subject to seasonal flooding. These areas are of recent origin, being formed largely from the sediment moved by the water in time of flood. As a consequence these deposits vary greatly from place to place. The top soil varies from loam to silt loam, with some areas being more sandy. In any one area rapid changes from loam, to sandy loam, to silty loam or silty clay loam can be expected. Stratification can be expected in some places. Within the valley of the Whitewater River the substratum ranges from silty clay to gravelly material with in short distances.

### C) EOLIAN DEPOSITS

Loess deposits are the only eolian or wind deposited materials occurring as mappable units in Fayette County.

### Loess Deposits

Most of the Wisconsin ground moraine is covered with a thin layer of wind-blown silt. In a few locations this silt deposit attains a considerable thickness and these areas have accordingly been mapped as loess deposits. Three feet of wind-blown silt was taken as the lower limit of such loess deposits; this depth coincides with the separation between the Birkbeck-Reesville-Manlove soil series and the Pincastle-Xenia-Russell soil series.

These silt deposits range from three to five feet thick in Fayette County and are underlain by Wisconsin till. The A-horizon is about one foot thick, and is silty loam. The B-horizon and C-horizons are also silty loams whose thicknesses vary according to the total depth of the loess deposit. The base of the loess deposit is marked by an abrupt change to a clay or silty clay (the Wisconsin till).

#### D) MISCELLANEOUS DEPOSITS

Several areas of highly organic topsoils have developed in depressions and poorly drained areas in the upland areas of Fayette County. There are also several areas in the southern part of the county where bedrock is at or near the surface. These regions are grouped under miscellaneous deposits.

##### Highly Organic Topsoil

Highly organic topsoil is found in depressions or other areas where drainage is retarded. Such areas are most commonly found in the Wisconsin ground and ridge moraines in the western part of the county. This area is relatively distant from the West Fork Whitewater River Valley and as a consequence stream erosion has not yet increased the local relief so that the natural drainage is slow. In general, these deposits are not deep in Fayette County and thus should not pose major construction problems. However these areas should be carefully field checked to determine the amount of material to be removed during construction.

##### Bedrock Near Surface

In several local areas in southern Fayette County bedrock is found at the surface or at very shallow depths. The bedrock consists of interbedded limestones and shales. The limestone tends to be thinly bedded and thus erosion often produces talus slopes of slabby limestone intermixed with shale fragments.

The most extensive areas of bedrock outcrop are found along Bear Creek and along the east side of the southern most portion of the West Fork Whitewater River valley in Fayette County, southeast of Alpine. Other important outcrops are found along the lower portions of Garrison Creek west and south-west of Alpine. However bedrock is encountered in most of



the lower portions of the small creeks in the southern part of the county.

The channels of these creeks are covered with small blocks of flaggy limestone.


## REFERENCES

1. Fenneman, N. M., "Physiography of the Eastern United States," McGraw-Hill, New York, 1938.
2. Gooding, A. M., "Pleistocene Terraces in the Upper Whitewater Drainage Basin, Southeastern Indiana," Earlham College Sci. Bull. 2, 65 p., 1957.
3. Gooding, A. M., "Illinoian and Wisconsin History in Southeastern Indiana," Geol. Soc. America, Field Trip Guidebook, Cincinnati Meeting 1961, pp. 99-128.
4. Gooding, A. M., "Illinoian and Wisconsin Glaciations in the Whitewater Basin, Southeastern Indiana, and Adjacent Areas," Journal of Geology Vol. 71, No. 6, pp. 665-682, 1963.
5. Malott, C. A., "The Physiography of Indiana," Handbook of Indiana Geology, Ind. Dept. Conservation, Pub. No. 21, 1922.
6. Parvis, M., "Airphoto Interpretation of Drainage Features of Fayette County," Joint Highway Research Project, Purdue University, Lafayette, Indiana, 1947.
7. "Soil Survey of Fayette and Union Counties, Indiana," United States Dept. of Agriculture, Soil Conservation Service and Purdue University Agricultural Experiment Station, Lafayette, Indiana, 1960.
8. "United States Census of Population, 1960," U. S. Bureau of Census, Government Printing Office, Washington, D. C., 1963.
9. Wayne, W. J., "Pleistocene Formations in Indiana," Indiana Dept. Conservation, Geological Survey Bulletin 25, Bloomington, Indiana, 1963.
10. Woods, K. B., "Highway Engineering Handbook," Section 9:- Distribution of Soils in North America, McGraw-Hill, 1960.
11. Yoder, E. J., "Principles of Pavement Design," John Wiley and Sons, Inc., New York, 1959, p. 569.


## GENERAL SOIL PROFILES

## GLACIAL DEPOSITS

## WISCONSIN CRAFT



## HLLINGMAN CRAFT



## FLUVIAL DEPOSITS

## TERRACES




## FLOOD PLAIN



## EOLIAN DEPOSITS



### MISCELLANEOUS DEPOSITS



ENGINEERING SOILS MAP  
FAYETTE COUNTY  
INDIANA

PREPARED FROM  
940 AIR AERIAL PHOTOGRAPHS  
BY  
JOINT HIGHSEA RESEARCH PROJECT, PURDUE UNIVERSITY

1990  
 1991  
 1992  
 1993  
 1994  
 1995  
 1996  
 1997  
 1998  
 1999  
 2000  
 2001  
 2002  
 2003  
 2004  
 2005  
 2006  
 2007  
 2008  
 2009  
 2010  
 2011  
 2012  
 2013  
 2014  
 2015  
 2016  
 2017  
 2018  
 2019  
 2020  
 2021  
 2022  
 2023  
 2024  
 2025  
 2026  
 2027  
 2028  
 2029  
 2030  
 2031  
 2032  
 2033  
 2034  
 2035  
 2036  
 2037  
 2038  
 2039  
 2040  
 2041  
 2042  
 2043  
 2044  
 2045  
 2046  
 2047  
 2048  
 2049  
 2050  
 2051  
 2052  
 2053  
 2054  
 2055  
 2056  
 2057  
 2058  
 2059  
 2060  
 2061  
 2062  
 2063  
 2064  
 2065  
 2066  
 2067  
 2068  
 2069  
 2070  
 2071  
 2072  
 2073  
 2074  
 2075  
 2076  
 2077  
 2078  
 2079  
 2080  
 2081  
 2082  
 2083  
 2084  
 2085  
 2086  
 2087  
 2088  
 2089  
 2090  
 2091  
 2092  
 2093  
 2094  
 2095  
 2096  
 2097  
 2098  
 2099  
 2100  
 2101  
 2102  
 2103  
 2104  
 2105  
 2106  
 2107  
 2108  
 2109  
 2110  
 2111  
 2112  
 2113  
 2114  
 2115  
 2116  
 2117  
 2118  
 2119  
 2120  
 2121  
 2122  
 2123  
 2124  
 2125  
 2126  
 2127  
 2128  
 2129  
 2130  
 2131  
 2132  
 2133  
 2134  
 2135  
 2136  
 2137  
 2138  
 2139  
 2140  
 2141  
 2142  
 2143  
 2144  
 2145  
 2146  
 2147  
 2148  
 2149  
 2150  
 2151  
 2152  
 2153  
 2154  
 2155  
 2156  
 2157  
 2158  
 2159  
 2160  
 2161  
 2162  
 2163  
 2164  
 2165  
 2166  
 2167  
 2168  
 2169  
 2170  
 2171  
 2172  
 2173  
 2174  
 2175  
 2176  
 2177  
 2178  
 2179  
 2180  
 2181  
 2182  
 2183  
 2184  
 2185  
 2186  
 2187  
 2188  
 2189  
 2190  
 2191  
 2192  
 2193  
 2194  
 2195  
 2196  
 2197  
 2198  
 2199  
 2200  
 2201  
 2202  
 2203  
 2204  
 2205  
 2206  
 2207  
 2208  
 2209  
 2210  
 2211  
 2212  
 2213  
 2214  
 2215  
 2216  
 2217  
 2218  
 2219  
 2220  
 2221  
 2222  
 2223  
 2224  
 2225  
 2226  
 2227  
 2228  
 2229  
 2230  
 2231  
 2232  
 2233  
 2234  
 2235  
 2236  
 2237  
 2238  
 2239  
 2240  
 2241  
 2242  
 2243  
 2244  
 2245  
 2246  
 2247  
 2248  
 2249  
 2250  
 2251  
 2252  
 2253  
 2254  
 2255  
 2256  
 2257  
 2258  
 2259  
 2260  
 2261  
 2262  
 2263  
 2264  
 2265  
 2266  
 2267  
 2268  
 2269  
 2270  
 2271  
 2272  
 2273  
 2274  
 2275  
 2276  
 2277  
 2278  
 2279  
 2280  
 2281  
 2282  
 2283  
 2284  
 2285  
 2286  
 2287  
 2288  
 2289  
 2290  
 2291  
 2292  
 2293  
 2294  
 2295  
 2296  
 2297  
 2298  
 2299  
 2300  
 2301  
 2302  
 2303  
 2304  
 2305  
 2306  
 2307  
 2308  
 2309  
 2310  
 2311  
 2312  
 2313  
 2314  
 2315  
 2316  
 2317  
 2318  
 2319  
 2320  
 2321  
 2322  
 2323  
 2324  
 2325  
 2326  
 2327  
 2328  
 2329  
 2330  
 2331  
 2332  
 2333  
 2334  
 2335  
 2336  
 2337  
 2338  
 2339  
 2340  
 2341  
 2342  
 2343  
 2344  
 2345  
 2346  
 2347  
 2348  
 2349  
 2350  
 2351  
 2352  
 2353  
 2354  
 2355  
 2356  
 2357  
 2358  
 2359  
 2360  
 2361  
 2362  
 2363  
 2364  
 2365  
 2366  
 2367  
 2368  
 2369  
 2370  
 2371  
 2372  
 2373  
 2374  
 2375  
 2376  
 2377  
 2378  
 2379  
 2380  
 2381  
 2382  
 2383  
 2384  
 2385  
 2386  
 2387  
 2388  
 2389  
 2390  
 2391  
 2392  
 2393  
 2394  
 2395  
 2396  
 2397  
 2398  
 2399  
 2400  
 2401  
 2402  
 2403  
 2404  
 2405  
 2406  
 2407  
 2408  
 2409  
 2410  
 2411  
 2412  
 2413  
 2414  
 2415  
 2416  
 2417  
 2418  
 2419  
 2420  
 2421  
 2422  
 2423  
 2424  
 2425  
 2426  
 2427  
 2428  
 2429  
 2430  
 2431  
 2432  
 2433  
 2434  
 2435  
 2436  
 2437  
 2438  
 2439  
 2440  
 2441  
 2442  
 2443  
 2444

194.7

00000000000000000000

# LEGEND

## PARENT MATERIALS

- (CLASSIFIED ACCORDING TO  
LAND FORM AND TOPOGRAPHY)
-  WESTERN COLORED MERGEL
  -  EASTERN COLORED MERGEL
  -  BLUE MERGEL
  -  TILLABLE
  -  PAVED ROAD
  -  UNPAVED ROAD
  -  GRAVEL

## MISCELLANEOUS

-  SMALL WET
-  SMALL SHADED TOPOGRAPHY
-  SAND AND GRAVEL
-  MARSH

## TEXTURAL SYMBOLS

- (BY COLOR AND TEXTURE  
SYMBOLS TO SHOW RELATIVE COMPOSITION)
-  GRAVEL
  -  SAND
  -  SILT
  -  CLAY

## TEXTURAL SYMBOLS FOR SOIL PROFILES

-  GRAVEL
-  SAND
-  SILT
-  CLAY
-  ORGANIC MATTER
-  UNSTABLE
-  STABLE

## ENGINEERING SOILS MAP FAYETTE COUNTY INDIANA

PREPARED FROM  
AERIAL PHOTOGRAPHS  
BY  
JOINT HIGHWAY RESEARCH PROJECT, PURDUE UNIVERSITY

PREPARED BY  
ALBERT E. HARRIS, CHAIRMAN  
SOIL CONSERVATION SERVICE  
AND  
BUREAU OF PUBLIC WORKS  
1947